

INFORMATION FOR PREVENTIVE VACCINATION, IN 2012

1. This Health Center is scheduled to conduct preventive vaccination for the students and the teaching staffs in 2012 in order to maintain and enhance their health.
2. Any students and teaching staffs including university personnel who wish to receive vaccination service is requested to take the following procedure.
 - a) **Preventive vaccination items** : uterine cervical cancer, A-type hepatitis ,B-type hepatitis, MMR(Measles, Mumps, Rubella), chicken pox

b) Date/time of scheduled vaccination

	Cervical cancer (vaccination 3times)	A-type hepatitis (vaccination 2times)	B-type hepatitis (vaccination 3times)	MMR (vaccination 2times)	Chicken pox (vaccination 2times)
Antibodies Test period	-	1pm ~ 4pm April 2(Mon)~6(Fri)		-	-
1 st vaccination schedule	1pm~4pm April 16(Mon)~18(Wed)				
2 nd vaccination schedule	1pm~4pm June 11(Mon)~15(Fri)	-	1pm~4pm May 14(Mon)~18(Fri)		

3 rd vaccination schedule	On September	-	-
--	--------------	---	---

- Be sure to confirm Attachment File for instruction before application.
- c) **Antibody Test & Vaccination place** : Industrial Medical Center, 1st basement level of the University Hospital
- d) **Subject of vaccination** : teaching staffs and students(& graduates) of this university
- e) **Antibody Test & Vaccination expense** (Card and cash payment acceptable)

Cost (one time))	Cervical cancer	A-type hepatitis	B-type hepatitis	MMA	Chicken pox
Antibody Test	-	KRW14,120	KRW12,610	-	-
Vaccination	KRW120,000	KRW50,000	KRW10,400	KRW13,450	KRW32,630

- To be paid on the date of vaccination
- f) **Others** : Dual vaccination permitted(Based on the guideline of Korea Centers for Disease Control & Prevention)
- g) **Supervision** : Infirmary (☎219-1596)
 - For further details, see each preventive vaccination-related data.(Homepage)

INSTRUCTION FOR VACCINATION

1. A person who should avoid vaccination

- A person with fever(due to flu & fatigue)
- A person with fever should be notified in advance..
- Acute respiratory patient
- A person allergic to chicken, egg & etc.
- A person with severe malnutrition
- Pregnant woman and female under menacme
- A person with cardiovascular disorder or hepatitis which is under acute or active stage
- A person with severe neurological disorder

2. Any side effect to be caused after vaccination

- Local stroke, swelling, pain or general fever, chills, headache, ennui, vomiting may be taken place but these symptoms are usually disappeared within 2-3 days.

3. Instruction after vaccination

- Please refrain from drinking and bathing on the date of vaccination. (Light shower is allowed.)